

CHIPPEWA VALLEY  
ORTHOPEDICS AND  
SPORTS MEDICINE

www.cvosm.com

**NATHANIEL J. STEWART, MD**  
Hip & Knee Expert

1200 OAKLEAF WAY STE A  
ALTOONA WI 54720  
TEL 715.832.1400

757 LAKELAND DR. STE B  
CHIPPEWA FALLS WI 54729  
TEL 715.723.8514

## DISCHARGE INSTRUCTIONS AFTER TOTAL HIP ARTHROPLASTY – POSTERIOR APPROACH

These instructions are meant to complement the information given by the nursing staff and physical therapists. They cover the most common questions and many of the areas that are unique to your surgical procedure.

### Wound Care

- Dressings are to be kept clean and dry. A small amount of clear drainage or bleeding may take place. When this is happening, the dressing should be changed daily.
- You may get the incision wet when showering five days after surgery. The shower should be brief (five minutes or so) and the wound patted dry with a clean towel.
- If you have staples or stitches, they are usually removed two weeks after surgery.
- If purulent material (thick white or greenish in color) is coming from the wound, or the wound is quite red on the edges, or you are having a temperature of 101.5 or higher, you should call my office and talk to me or the doctor on call.

### Weight Bearing

- Unless the physical therapist has told you otherwise, you can put as much weight on your hip as feels comfortable. Depending on your age, strength, and coordination, most people use the walker or crutches for three weeks with a gradual progression to a crutch or cane over the next few weeks.

### Avoiding Dislocation

- Physical therapy in the hospital after surgery will review with you this important topic. The restrictions are not to bend the hip beyond 90 degrees, not to cross your legs, and not to reach down for the outside part of your shoe. The 90 degree rule is most often broken by some one sitting on a surface that is too low.

### Outpatient Physical Therapy


- The most important goal of physical therapy is to avoid dislocation. The next goal is to restore normal gait (walking pattern) and to improve strength and flexibility, returning you to your normal activities. Speak with your physical therapist regarding your goals.

### Swelling

- Swelling to some degree is common after joint replacement. To reduce swelling, elevation is very helpful. Lying down, with your “toes above your nose” at least 30 minutes of every 2 hours is a good initial recommendation- you may need more or less time. The white stockings are also designed to reduce swelling. Excessive swelling with pain or tenderness in the calf, redness of the lower leg, and/or increased warmth of the leg, can be a sign of a blood clot. Patients frequently call to ask if their swelling is too much, and this is impossible to tell over the phone. If you feel you are excessively swollen with pain or tenderness, it is best investigated by exam and may require an ultrasound to rule out a blood clot.

### Cold therapy

- Cold therapy can help reduce pain and swelling. It should be used intermittently for the first few weeks, as long as you are having some swelling or discomfort in the area of the surgery. Typically, patients use it two or three times a day for up to 30-minutes at a time. It should not cause pain and there should be a thin



www.cvosm.com

**NATHANIEL J. STEWART, MD**  
Hip & Knee Expert

1200 OAKLEAF WAY STE A  
ALTOONA WI 54720  
TEL 715.832.1400

757 LAKELAND DR. STE B  
CHIPPEWA FALLS WI 54729  
TEL 715.723.8514

towel between your surgical site and the cooling device. Cooling devices can be commercial devices that pump cold water around your joint or a frozen gel pack. In any case, it should feel better with the cold applied and reduce your discomfort in the first few months after surgery.

### **Driving**

- To drive you must no longer be taking pain pills (that is narcotics; Tylenol and anti-inflammatory medications are allowed). Also, you must feel strong and alert. For most people this is between 1-2 weeks after surgery, depending on the side of your surgery.

### **Preventing Blood Clots**

- The first line of defense is moving your ankles up and down. This action circulates the blood from your legs back to your heart, preventing a clot.
- Patients are put on either aspirin or Coumadin postoperatively to reduce the risk of developing a blood clot. We select the choice of medication for blood clot prevention very carefully using a list of risk factor criteria that would increase an individual's risk of developing a blood clot. Patients that are at a high risk for developing a blood clot are put on Coumadin for a total of three weeks. Patients respond very differently to the same dose of Coumadin, and so the effects of the drug need to be monitored with a blood test called Protime. Initially your Protime will be checked at a laboratory near you on Monday and Thursday mornings, and the results called o Dr. Stewart's office. You will be contacted if you need to adjust your dose
- Compressive Stockings: These can prevent blood from pooling in your leg if you can use them correctly. You don't need to sleep with them on. You put them on first thing in the morning before your leg can swell. They are recommended for the first three weeks after surgery. If they just keep curling up and binding in, then they are not working and should be discontinued.


### **Sleeping**

Surgery can throw off your normal sleep wake cycle, leaving you staring at the wall at 3 AM. To reset your sleep wake cycle, follow this program:

- No naps during the day
- Decrease your narcotic use
- Only one caffeinated beverage per day and it must be before 10AM
- No exercise or excitement after 6PM
- Go to bed and get up at the same time every day no matter if you are sleepy or not
- Allow slightly less total time for sleep than you would normally
- While sleeping on your back, use the abductor pillow from the hospital. At 4 weeks after surgery, you can roll to your opposite side with pillows in between your knees.

### **Narcotic Pain Medication**

- You will need narcotic pain medication after you leave the hospital. They can be addictive, and you will be weaned from them as tolerated. If you are running low, please call the office during regular business hours for a refill. In general, refills will not be made by the doctor on call after hours, so plan ahead



www.cvosm.com

**NATHANIEL J. STEWART, MD**  
Hip & Knee Expert

1200 OAKLEAF WAY STE A  
**ALTOONA** WI 54720  
TEL 715.832.1400

757 LAKELAND DR. STE B  
**CHIPPEWA FALLS** WI 54729  
TEL 715.723.8514

### **Sexual Relations**

- Wait until your general health has recovered, often 4 to 6 weeks. Avoid positions that would put your hip at risk of dislocation

### **Dental Work (or other invasive procedures)**

- Unless it is an emergency, it is best to avoid dental work or other invasive procedures for 3 months after a joint replacement. When undergoing an invasive procedure, we recommend antibiotics to prevent bacterial from getting in the blood stream and infecting the joint. Please call our office for an antibiotic prescription prior to any dental work.

### **Phone Calls**

- If you have questions or problems please call 715-832-1400. Some questions can be answered over the phone, others cannot. You may need to be seen sooner than expected to check out your symptoms.

### **Follow Up**

- Make sure an appointment has been scheduled for you at Chippewa Valley Orthopedics & Sports Medicine for approximately 10-14 days after surgery.

**Nathaniel Stewart, M.D.**  
**Chippewa Valley Orthopedics & Sports Medicine**  
**(715) 832-1400 or (715) 723-8514, or (800) 322-1747**